

Product name:	IKK beta Rabbit Monoclonal Antibody
Cat number:	MABN85700
Conjugate:	Unconjugated
Size:	100µL
Clone:	Monoclonal
Concentration:	1mg/ml
Host:	Rabbit
Isotype:	IgG
Immunogen:	A synthetic peptide of human IKK beta
Reactivity:	Human, Mouse, Rat
Applications:	WB 1:500-1:1000, ICC 1:50-1:200
Molecular Weight:	Calculated MW: 87 kDa; Observed MW: 87 kDa
Purification:	Affinity Purification
Form:	Liquid
Buffer:	Purified antibody in TBS with 0.05% sodium azide, 0.05% BSA and 50% glycerol.
Storage:	Store at 4°C short term. Aliquot and store at -20°C for 12 months. Avoid freeze/thaw cycles.
Background:	The NF-κB/Rel transcription factors are present in the cytosol in an inactive state, complexed with the inhibitory IκB proteins (1-3). Most agents that activate NF-κB do so through a common pathway based on phosphorylation-induced, proteasome-mediated degradation of IκB (3-7). The key regulatory step in this pathway involves activation of a high molecular weight IκB kinase (IKK) complex whose catalysis is generally carried out by three tightly associated IKK subunits.