

Product name:	JIP-3 Rabbit Polyclonal Antibody
Cat number:	ABN12837
Conjugate:	Unconjugated
Size:	100µL
Clone:	Polyclonal
Concentration:	1mg/ml
Host:	Rabbit
Isotype:	IgG
Immunogen:	The antiserum was produced against synthesized peptide derived from human JIP3. AA range:621-670
Reactivity:	Human,Mouse
Applications:	WB 1:500-1:2000,IHC 1:100-1:300,ICC/IF 1:200-1:1000,ELISA 1:10000-1:20000
Molecular Weight:	170kDa
Purification:	Affinity purification
Form:	Liquid
Buffer:	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.
Storage:	Store at 4°C short term. Aliquot and store at -20°C for 12 months. Avoid freeze/thaw cycles.

Background:

The protein encoded by this gene shares similarity with the product of *Drosophila* *syd* gene, required for the functional interaction of kinesin I with axonal cargo. Studies of the similar gene in mouse suggested that this protein may interact with, and regulate the activity of numerous protein kinases of the JNK signaling pathway, and thus function as a scaffold protein in neuronal cells. The *C. elegans* counterpart of this gene is found to regulate synaptic vesicle transport possibly by integrating JNK signaling and kinesin-1 transport. Several alternatively spliced transcript variants of this gene have been described, but the full-length nature of some of these variants has not been determined. [provided by RefSeq, Jul 2008],function:The JNK-interacting protein (JIP) group of scaffold proteins selectively mediates JNK signaling by aggregating specific components of the MAPK cascade to form a functional JNK signaling module. May function as a regulator of vesicle transport, through interactions with the JNK-signaling components and motor proteins.,PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Belongs to the JIP scaffold family.,subunit:Forms homo- or heterooligomeric complexes. The central region of MAPK8IP3 interacts with the C-terminal of MAPK8IP2 but not MAPK8IP1. Binds specific components of the JNK signaling pathway namely MAPK8, MAPK9 and MAPK10 to the N-terminal region, MAP2K4 and MAP2K7 to the central region and MAP3K11 to the C-terminal region. Binds the TPR motif-containing C-terminal of kinesin light chain, KLC1. Pre-assembled MAPK8IP1 scaffolding complexes are then transported as a cargo of kinesin, to the required subcellular location.,