




Product name:	[KD Validated] COX5A Rabbit mAb
Cat number:	AB25751
Conjugate:	Unconjugated
Size:	100 ug
Clone:	ARC67578
Concentration:	1mg/ml
Host:	Rabbit
Isotype:	IgG
Immunogen:	Recombinant fusion protein containing a sequence corresponding to amino acids 1-150 of human COX5A.
Reactivity:	Human,Mouse,Rat
Applications:	WB 1:2000 - 1:8000 IHC-P 1:300 - 1:3000 IF/ICC 1:200 - 1:800 ELISA Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.
Molecular Weight:	13kDa
Purification:	Affinity purification
Form:	liquid
Buffer:	PBS with 0.05% proclin300,0.05% BSA,50% glycerol,pH7.3.
Storage:	Store at -20°C. Avoid freeze / thaw cycles.
Background:	Cytochrome c oxidase (COX) is the terminal enzyme of the mitochondrial respiratory chain. It is a multi-subunit enzyme complex that couples the transfer of electrons from cytochrome c to molecular oxygen and contributes to a proton electrochemical gradient across the inner mitochondrial membrane. The complex consists of 13 mitochondrial- and nuclear-encoded subunits. The mitochondrially-encoded subunits perform the electron transfer of proton pumping activities. The functions of the nuclear-encoded subunits are unknown but they may play a role in the regulation and assembly of the complex. This gene encodes the nuclear-encoded subunit Va of the human mitochondrial respiratory chain enzyme. A pseudogene COX5AP1 has been found in chromosome 14q22.




Western blot analysis of lysates from wild type (WT) and COX5A knockdown (KD) HeLa cells using [KD Validated] COX5A Rabbit mAb (A25751) at 1:2000 dilution incubated overnight at 4°C. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25 µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020) Exposure time: 30 s.




Western blot analysis of lysates from 293T cells using [KD Validated] COX5A Rabbit mAb (A25751) at 1:2000 dilution incubated overnight at 4°C. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25 µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020) Exposure time: 30 s.




Immunohistochemistry analysis of paraffin-embedded Human liver tissue using [KD Validated] COX5A Rabbit mAb (A25751) at a dilution of 1:300 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate Bufferr (pH 6.0) prior to IHC staining.




Immunohistochemistry analysis of paraffin-embedded Human kidney tissue using [KD Validated] COX5A Rabbit mAb (A25751) at a dilution of 1:300 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate Bufferr (pH 6.0) prior to IHC staining.




Immunohistochemistry analysis of paraffin-embedded Human colon tissue using [KD Validated] COX5A Rabbit mAb (A25751) at a dilution of 1:300 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate Bufferr (pH 6.0) prior to IHC staining.




Confocal imaging of HeLa cells using [KD Validated] COX5A Rabbit mAb (A25751, dilution 1:200) followed by a further incubation with Cy3 Goat Anti-Rabbit IgG (H+L) (AS007, dilution 1:500) (Red). The cells were counterstained with α-Tubulin Mouse mAb (A0212, dilution 1:400) followed by incubation with AbRio® 488-conjugated Goat Anti-Mouse IgG (H+L) Ab (AS076, dilution 1:500) (Green). DAPI was used for nuclear staining (Blue). Objective: 100x.



Confocal imaging of 293T cells using [KD Validated] COX5A Rabbit mAb (A25751, dilution 1:200) followed by a further incubation with Cy3 Goat Anti-Rabbit IgG (H+L) (A5007, dilution 1:500) (Red). DAPI was used for nuclear staining (Blue). Objective: 100x.



Confocal imaging of 293T cells using [KD Validated] COX5A Rabbit mAb (A25751, dilution 1:200) followed by a further incubation with Cy3 Goat Anti-Rabbit IgG (H+L) (A5007, dilution 1:500) (Red). DAPI was used for nuclear staining (Blue). High pressure antigen retrieval performed with 0.01M Citrate Buffer (pH 6.0) prior to IF staining. Objective: 40x.



Confocal imaging of paraffin-embedded Rat brain tissue using [KD Validated] COX5A Rabbit mAb (A25751, dilution 1:200) followed by a further incubation with Cy3 Goat Anti-Rabbit IgG (H+L) (A5007, dilution 1:500) (Red). DAPI was used for nuclear staining (Blue). Microwave antigen retrieval performed with 0.01M Citrate Buffer (pH 6.0) prior to IF staining. Objective: 40x.