

|                          |                                                                                                     |
|--------------------------|-----------------------------------------------------------------------------------------------------|
| <b>Product name:</b>     | KIF4A Rabbit Polyclonal Antibody                                                                    |
| <b>Cat number:</b>       | ABN13015                                                                                            |
| <b>Conjugate:</b>        | Unconjugated                                                                                        |
| <b>Size:</b>             | 100µL                                                                                               |
| <b>Clone:</b>            | Polyclonal                                                                                          |
| <b>Concentration:</b>    | 1mg/ml                                                                                              |
| <b>Host:</b>             | Rabbit                                                                                              |
| <b>Isotype:</b>          | IgG                                                                                                 |
| <b>Immunogen:</b>        | The antiserum was produced against synthesized peptide derived from human KIF4A. AA range:1171-1220 |
| <b>Reactivity:</b>       | Human,Rat,Mouse                                                                                     |
| <b>Applications:</b>     | WB 1:500-1:2000,IHC 1:50-1:300,ELISA 1:2000-1:20000                                                 |
| <b>Molecular Weight:</b> | 140kDa                                                                                              |
| <b>Purification:</b>     | Affinity purification                                                                               |
| <b>Form:</b>             | Liquid                                                                                              |
| <b>Buffer:</b>           | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.                  |
| <b>Storage:</b>          | Store at 4°C short term. Aliquot and store at -20°C for 12 months. Avoid freeze/thaw cycles.        |

**Background:**

kinesin family member 4A(KIF4A) Homo sapiens This gene encodes a member of the kinesin 4 subfamily of kinesin related proteins. The encoded protein is an ATP dependent microtubule-based motor protein that is involved in the intracellular transport of membranous organelles. This protein also associates with condensed chromosome arms and may be involved in maintaining chromosome integrity during mitosis. This protein may also be involved in the organization of the central spindle prior to cytokinesis. A pseudogene of this gene is found on chromosome X.[provided by RefSeq, Mar 2010],function:Motor protein that translocates PRC1 to the plus ends of interdigitating spindle microtubules during the metaphase to anaphase transition, an essential step for the formation of an organized central spindle midzone and midbody and for successful cytokinesis. May play a role in mitotic chromosomal positioning and bipolar spindle stabilization.,sequence caution:Contaminating sequence. Potential poly-A sequence.,similarity:Belongs to the kinesin-like protein family.,similarity:Belongs to the kinesin-like protein family. Chromokinesin subfamily.,similarity:Contains 1 kinesin-motor domain.,subcellular location:Not present in the nucleolus. In early mitosis, associated with the mitotic spindle, in anaphase, localized to the spindle midzone and, in telophase and cytokinesis, to the midbody. In late cytokinesis, found in the center of the midbody. Associated with chromosomes at all stages of mitosis.,subunit:Interacts with unphosphorylated PRC1 during late mitosis.,tissue specificity:Highly expressed in hematopoietic tissues, fetal liver, spleen, thymus and adult thymus and bone marrow. Lower levels are found in heart, testis, kidney, colon and lung.,tissue specificity:Specifically expressed in testis.,