

Product name:	Liprin β 1 Rabbit Polyclonal Antibody
Cat number:	ABN13329
Conjugate:	Unconjugated
Size:	100 μ L
Clone:	Polyclonal
Concentration:	1mg/ml
Host:	Rabbit
Isotype:	IgG
Immunogen:	The antiserum was produced against synthesized peptide derived from human LIPB1. AA range:300-349
Reactivity:	Human,Mouse
Applications:	WB 1:500-1:2000,IHC 1:50-1:300
Molecular Weight:	115kDa
Purification:	Affinity purification
Form:	Liquid
Buffer:	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.
Storage:	Store at 4°C short term. Aliquot and store at -20°C for 12 months. Avoid freeze/thaw cycles.

Background:

The protein encoded by this gene is a member of the LAR protein-tyrosine phosphatase-interacting protein (liprin) family. Liprins interact with members of LAR family of transmembrane protein tyrosine phosphatases, which are known to be important for axon guidance and mammary gland development. It has been proposed that liprins are multivalent proteins that form complex structures and act as scaffolds for the recruitment and anchoring of LAR family of tyrosine phosphatases. This protein was found to interact with S100A4, a calcium-binding protein related to tumor invasiveness and metastasis. In vitro experiment demonstrated that the interaction inhibited the phosphorylation of this protein by protein kinase C and protein kinase CK2. Alternatively spliced transcript variants encoding distinct isoforms have been reported. [provided by RefSeq, Jul 2008],domain:The N-terminal coiled coil regions mediate homodimerization preferentially and heterodimerization type beta/beta. The C-terminal, non-coiled coil regions mediate heterodimerization type beta/alpha and interaction with S100A4.,function:May regulate the disassembly of focal adhesions. Did not bind receptor-like tyrosine phosphatases type 2A.,similarity:Belongs to the liprin family. Liprin-beta subfamily.,similarity:Contains 3 SAM (sterile alpha motif) domains.,subunit:Forms homodimers and heterodimers. Interacts with S100A4 in a calcium-dependent mode.,tissue specificity:Widely expressed. Absent in liver.,