

Product name:	LPD lipase Rabbit Polyclonal Antibody
Cat number:	ABN13391
Conjugate:	Unconjugated
Size:	100µL
Clone:	Polyclonal
Concentration:	1mg/ml
Host:	Rabbit
Isotype:	IgG
Immunogen:	The antiserum was produced against synthesized peptide derived from human LIPI. AA range:289-338
Reactivity:	Human,Rat,Mouse
Applications:	WB 1:500-1:2000,IHC 1:100-1:300,ICC/IF 1:50-1:200,ELISA 1:5000-1:20000
Molecular Weight:	53kDa
Purification:	Affinity purification
Form:	Liquid
Buffer:	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.
Storage:	Store at 4°C short term. Aliquot and store at -20°C for 12 months. Avoid freeze/thaw cycles.
Background:	The protein encoded by this gene is a phospholipase that hydrolyzes phosphatidic acid to produce lysophosphatidic acid. Defects in this gene are a cause of susceptibility to familial hypertriglyceridemia. This gene is also expressed at high levels in Ewing family tumor cells. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Dec 2014],disease:Defects in LIPI may be a cause of susceptibility to familial hypertriglyceridemia [MIM:145750]. Familial hypertriglyceridemia is a common inherited disorder in which the concentration of very low density lipoprotein (VLDL) is elevated in the plasma. This leads to increased risk of heart disease, obesity, and pancreatitis.,enzyme regulation:Inhibited by sodium vanadate.,function:Hydrolyzes specifically phosphatidic acid (PA) to produce lysophosphatidic acid (LPA).,PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Belongs to the AB hydrolase superfamily. Lipase family.,subcellular location:May associate with lipid draft.,subunit:Interacts with heparin with a high affinity.,tissue specificity:Expressed in testis. Expressed exclusively at the connecting piece of the sperm.,