

Product name:	MAGI-2 Rabbit Polyclonal Antibody
Cat number:	ABN13599
Conjugate:	Unconjugated
Size:	100µL
Clone:	Polyclonal
Concentration:	1mg/ml
Host:	Rabbit
Isotype:	IgG
Immunogen:	The antiserum was produced against synthesized peptide derived from human MAGI2. AA range:221-270
Reactivity:	Human,Mouse,Rat
Applications:	WB 1:500-1:2000,IHC 1:100-1:300,ICC/IF 1:50-1:200,ELISA 1:10000-1:20000
Molecular Weight:	156kDa
Purification:	Affinity purification
Form:	Liquid
Buffer:	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.
Storage:	Store at 4°C short term. Aliquot and store at -20°C for 12 months. Avoid freeze/thaw cycles.

Background:

The protein encoded by this gene interacts with atrophin-1. Atrophin-1 contains a polyglutamine repeat, expansion of which is responsible for dentatorubral and pallidoluysian atrophy. This encoded protein is characterized by two WW domains, a guanylate kinase-like domain, and multiple PDZ domains. It has structural similarity to the membrane-associated guanylate kinase homologue (MAGUK) family. [provided by RefSeq, Jul 2008],function:Seems to act as scaffold molecule at synaptic junctions by assembling neurotransmitter receptors and cell adhesion proteins. May play a role in regulating activin-mediated signaling in neuronal cells. Enhances the ability of PTEN to suppress AKT1 activation.,similarity:Belongs to the MAGUK family.,similarity:Contains 1 guanylate kinase-like domain.,similarity:Contains 2 WW domains.,similarity:Contains 6 PDZ (DHR) domains.,subcellular location:Membrane-associated in synaptosomes.,subunit:Interacts via its WW domains with DRPLA. Interacts via its second PDZ domain with PTEN unphosphorylated C-terminus; this interaction diminishes the degradation rate of PTEN (By similarity). Interacts through its guanylate kinase domain with DLGAP1 (By similarity). Interacts through the PDZ domains with GRIN2A, GRID2 and NLGN1 (By similarity). Interacts with CTNND2, CTNNB1, MAGUIN-1, ACVR2A, SMAD2 and SMAD3 (By similarity). Part of a complex consisting of AIP1, ACVR2A, ACVR1B and SMAD3 (By similarity). May interact with HTR2A (By similarity). Interacts with IGSF9, RAPGEF2 and HTR4 (By similarity). Interacts with DDN.,tissue specificity:Specifically expressed in brain.,