

Product name:	MIP-1 α Rabbit Polyclonal Antibody
Cat number:	ABN13908
Conjugate:	Unconjugated
Size:	100 μ L
Clone:	Polyclonal
Concentration:	1mg/ml
Host:	Rabbit
Isotype:	IgG
Immunogen:	The antiserum was produced against synthesized peptide derived from human MIP-1 α . AA range:26-75
Reactivity:	Human,Rat,Mouse
Applications:	IHC 1:100-1:300,ICC/IF 1:50-1:200,ELISA 1:10000-1:20000
Purification:	Affinity purification
Form:	Liquid
Buffer:	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.
Storage:	Store at 4°C short term. Aliquot and store at -20°C for 12 months. Avoid freeze/thaw cycles.
Background:	This locus represents a small inducible cytokine. The encoded protein, also known as macrophage inflammatory protein 1 α, plays a role in inflammatory responses through binding to the receptors CCR1, CCR4 and CCR5. Polymorphisms at this locus may be associated with both resistance and susceptibility to infection by human immunodeficiency virus type 1.[provided by RefSeq, Sep 2010],function:Monokine with inflammatory and chemokinetic properties. Binds to CCR1, CCR4 and CCR5. One of the major HIV-suppressive factors produced by CD8+ T-cells. Recombinant MIP-1-α induces a dose-dependent inhibition of different strains of HIV-1, HIV-2, and simian immunodeficiency virus (SIV).,induction:By TPA or PHA (TPA = 12-O-tetradecanoyl phorbol-13 acetate (tumor promoter); PHA = phytohemagglutinin (T-cell mitogen)).,online information:Macrophage inflammatory protein entry,PTM:N-terminal processed form LD78-α(4-69) is produced by proteolytic cleavage after secretion from HTLV1-transformed T-cells.,similarity:Belongs to the intercrine beta (chemokine CC) family.,subunit:Self-associates. Also heterodimer of MIP-1-α(4-69) and MIP-1-β(3-69).,