

|                          |                                                                                                                                             |
|--------------------------|---------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Product name:</b>     | NFATc4 (phospho Ser676) Rabbit Polyclonal Antibody                                                                                          |
| <b>Cat number:</b>       | ABN05088                                                                                                                                    |
| <b>Conjugate:</b>        | Unconjugated                                                                                                                                |
| <b>Size:</b>             | 100µL                                                                                                                                       |
| <b>Clone:</b>            | Polyclonal                                                                                                                                  |
| <b>Concentration:</b>    | 1mg/ml                                                                                                                                      |
| <b>Host:</b>             | Rabbit                                                                                                                                      |
| <b>Isotype:</b>          | IgG                                                                                                                                         |
| <b>Immunogen:</b>        | The antiserum was produced against synthesized peptide derived from human NFAT3 around the phosphorylation site of Ser676. AA range:642-691 |
| <b>Reactivity:</b>       | Human,Mouse                                                                                                                                 |
| <b>Applications:</b>     | WB 1:500-1:2000,IHC 1:100-1:300,ICC/IF 1:50-1:200,ELISA 1:5000-1:10000                                                                      |
| <b>Molecular Weight:</b> | 120kDa                                                                                                                                      |
| <b>Purification:</b>     | Affinity purification                                                                                                                       |
| <b>Form:</b>             | Liquid                                                                                                                                      |
| <b>Buffer:</b>           | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.                                                          |
| <b>Storage:</b>          | Store at 4°C short term. Aliquot and store at -20°C for 12 months. Avoid freeze/thaw cycles.                                                |

**Background:**

This gene encodes a member of the nuclear factor of activated T cells (NFAT) protein family. The encoded protein is part of a DNA-binding transcription complex. This complex consists of at least two components: a preexisting cytosolic component that translocates to the nucleus upon T cell receptor stimulation and an inducible nuclear component. NFAT proteins are activated by the calmodulin-dependent phosphatase, calcineurin. The encoded protein plays a role in the inducible expression of cytokine genes in T cells, especially in the induction of interleukin-2 and interleukin-4. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Jan 2014], domain: Rel Similarity Domain (RSD) allows DNA-binding and cooperative interactions with AP1 factors., function: Plays a role in the inducible expression of cytokine genes in T-cells, especially in the induction of the IL-2 and IL-4. Transcriptionally repressed by estrogen receptors; this inhibition is further enhanced by estrogen. Increases the transcriptional activity of PPARG and has a direct role in adipocyte differentiation. May play an important role in myotube differentiation. May play a critical role in cardiac development and hypertrophy. May play a role in deafferentation-induced apoptosis of sensory neurons., PTM: Phosphorylated by NFATC-kinases; dephosphorylated by calcineurin. Phosphorylated on Ser-168 and Ser-170 by FRAP1, IRAK1, MAPK7 and MAPK14, on Ser-213 and Ser-217 by MAPK8 and MAPK9, and on Ser-289 and Ser-344 by RPS6KA3. Phosphorylated by GSK3B., PTM: Ubiquitinated, leading to its degradation by the proteasome and reduced transcriptional activity. Ubiquitination and reduction in transcriptional activity can be further facilitated through GSK3B-dependent phosphorylation. Polyubiquitin linkage is mainly through 'Lys-48'. , similarity: Contains 1 IPT/TIG domain., similarity: Contains 1 RHD (Rel-like) domain., subcellular location: Cyttoplasmic for the phosphorylated form and nuclear after activation that is controlled by calcineurin-mediated dephosphorylation. Rapid nuclear exit of NFATC is thought to be one mechanism by which cells distinguish between sustained and transient calcium signals. The subcellular localization of NFATC plays a key role in the regulation of gene transcription., subunit: Member of the multicomponent NFATC transcription complex that consists of at least two components, a pre-existing cytoplasmic component NFATC2 and an inducible nuclear component NFATC1. Other members such as NFATC4, NFATC3 or members of the activating protein-1 family, MAF, GATA4 and Cbp/p300 can also bind the complex. NFATC proteins bind to DNA as monomers. Interacts with CREBBP, GATA4, IRAK1, MAPK8, MAPK9 and RPS6KA3., tissue specificity: Highly expressed in placenta, lung, kidney, testis and ovary. Weakly expressed in spleen and thymus. Not expressed in peripheral blood lymphocytes. Detected in hippocampus.,