





Product name:	NFKB p65
Cat number:	AB-82033
Conjugate:	Unconjugated
Size:	100 ug
Clone:	POLY
Concentration:	1mg/ml
Host:	Rb
Isotype:	IgG
Immunogen:	Recombinant fusion protein containing a sequence corresponding to amino acids 50-180 of human NF-kB p65.
Reactivity:	Hu, Ms, Rt
Applications:	Western Blot: 1:1000 Immunohistochemistry: 1:50 - 1:100 Immunofluorescence: 1:50 - 1:100 Immunoprecipitation: 1:50 - 1:100
Molecular Weight:	65 kDa
Purification:	Aff. Pur.
Form:	Liquid
Buffer:	PBS with 0.02% sodium azide, 50% glycerol, pH7.3.
Storage:	Store at -20°C. Avoid freeze / thaw cycles.
Background:	NF-kappa-B is a ubiquitous transcription factor involved in several biological processes. It is held in the cytoplasm in an inactive state by specific inhibitors. Upon degradation of the inhibitor, NF-kappa-B moves to the nucleus and activates transcription of specific genes. NF-kappa-B is composed of NFKB1 or NFKB2 bound to either REL, RELA, or RELB. The most abundant form of NF-kappa-B is NFKB1 complexed with the product of this gene, RELA. Four transcript variants encoding different isoforms have been found for this gene.



Immunohistochemistry of paraffin-embedded rat lung using NF-kB p65 Antibody at dilution of 1:100 (40x lens).




Immunohistochemistry of paraffin-embedded rat kidney using NF-kB p65 Antibody at dilution of 1:100 (40x lens).



Immunofluorescence analysis of U2OS cells using NF-kB p65 antibody.

Fig.5C



蛋白上样量: 15μl;

抗体使用量: 8μl;

抗体稀释度: 1:500;

实验组织: ox-LDL刺激的Raw264.7细胞

Application: Western blot (WB) Sample: RAW264.7 Loading Amount: 15 ug Dilution: 1:500
Incubation time: over night Temperature 4 °C Exposure: 15 seconds Observed Band: 65 kDa
Remarks Western blot analysis of extracts of mouse atherosclerosis arterial tissue, using
RELA antibody at 1:500 dilution. The band is specific and clear.