

Product name:	O-FucT-2 Rabbit Polyclonal Antibody
Cat number:	ABN15117
Conjugate:	Unconjugated
Size:	100µL
Clone:	Polyclonal
Concentration:	1mg/ml
Host:	Rabbit
Isotype:	IgG
Immunogen:	The antiserum was produced against synthesized peptide derived from human POFUT2. AA range:361-410
Reactivity:	Human,Mouse
Applications:	IHC 1:100-1:300,ICC/IF 1:50-1:200,ELISA 1:10000-1:20000
Purification:	Affinity purification
Form:	Liquid
Buffer:	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.
Storage:	Store at 4°C short term. Aliquot and store at -20°C for 12 months. Avoid freeze/thaw cycles.
Background:	Fucose is typically found as a terminal modification of branched chain glycoconjugates, but it also exists in direct O-linkage to serine or threonine residues within cystine knot motifs in epidermal growth factor (EGF; MIM 131530)-like repeats or thrombospondin (THBS; see MIM 188060) type-1 repeats. POFUT2 is an O-fucosyltransferase that use THBS type-1 repeats as substrates (Luo et al., 2006 [PubMed 16464857]).[supplied by OMIM, Mar 2008],catalytic activity:Transfers an alpha-L-fucosyl residue from GDP-beta-L-fucose to the serine hydroxy group of a protein acceptor.,function:Catalyzes the reaction that attaches fucose through an O-glycosidic linkage to a conserved serine or threonine residue in thrombospondin type 1 repeats.,online information:GlycoGene database,online information:Peptide-O-fucosyltransferase 2,pathway:Protein modification; protein glycosylation.,similarity:Belongs to the glycosyltransferase 68 family.,tissue specificity:Isoform A is expressed in fetal liver and peripheral blood lymphocytes. Isoform B is expressed in spleen, lung, testis, bone marrow, thymus, pancreas, prostate, fetal brain, fetal liver and fetal kidney. Isoform C is expressed in brain, heart, spleen, liver, lung, stomach, testis, placenta, skin, thymus, pancreas, mammary gland, prostate, fetal brain, fetal liver and fetal heart.,