

Product name:	OASL Rabbit Polyclonal Antibody
Cat number:	ABN15079
Conjugate:	Unconjugated
Size:	100µL
Clone:	Polyclonal
Concentration:	1mg/ml
Host:	Rabbit
Isotype:	IgG
Immunogen:	Synthesized peptide derived from human OASL. at AA range: 1-50
Reactivity:	Rat,Mouse
Applications:	WB 1:500-1:2000,ELISA 1:10000-1:20000
Molecular Weight:	60kDa
Purification:	Affinity purification
Form:	Liquid
Buffer:	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.
Storage:	Store at 4°C short term. Aliquot and store at -20°C for 12 months. Avoid freeze/thaw cycles.
Background:	caution:This may not be the true ortholog of mouse OASL.,function:Does not have 2'-5'-OAS activity, but binds double-stranded RNA and DNA.,induction:By interferons.,similarity:Belongs to the 2-5A synthetase family.,similarity:Contains 2 ubiquitin-like domains.,subunit:Specifically interacts with the ligand binding domain of the thyroid receptor (TR). TRIP14 does not require the presence of thyroid hormone for its interaction. Binds MBD1.,tissue specificity:Expressed in most tissues, with the highest levels in primary blood Leukocytes and other hematopoietic system tissues, colon, stomach and to some extent in testis.,caution:This may not be the true ortholog of mouse OASL.,function:Does not have 2'-5'-OAS activity, but binds double-stranded RNA and DNA.,induction:By interferons.,similarity:Belongs to the 2-5A synthetase family.,similarity:Contains 2 ubiquitin-like domains.,subunit:Specifically interacts with the ligand binding domain of the thyroid receptor (TR). TRIP14 does not require the presence of thyroid hormone for its interaction. Binds MBD1.,tissue specificity:Expressed in most tissues, with the highest levels in primary blood Leukocytes and other hematopoietic system tissues, colon, stomach and to some extent in testis.,