



Product name:	p21 Polyclonal Antibody
Cat number:	AB-83710
Conjugate:	Unconjugated
Size:	100ug
Clone:	Polyclonal
Concentration:	1 mg/mL
Host:	Rabbit
Isotype:	IgG
Immunogen:	Recombinant fusion protein of human p21 (NP_000380.1).
Reactivity:	Human;Mouse;Rat
Applications:	WB 1:500-1:2000;IF 1:50-1:200
Molecular Weight:	21 kDa
Purification:	Affinity purification
Buffer:	Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol.
Storage:	Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.
Background:	This gene encodes a potent cyclin-dependent kinase inhibitor. The encoded protein binds to and inhibits the activity of cyclin-cyclin-dependent kinase2 or -cyclin-dependent kinase4 complexes, and thus functions as a regulator of cell cycle progression at G1. The expression of this gene is tightly controlled by the tumor suppressor protein p53, through which this protein mediates the p53-dependent cell cycle G1 phase arrest in response to a variety of stress stimuli. This protein can interact with proliferating cell nuclear antigen, a DNA polymerase accessory factor, and plays a regulatory role in S phase DNA replication and DNA damage repair. This protein was reported to be specifically cleaved by CASP3-like caspases, which thus leads to a dramatic activation of cyclin-dependent kinase2, and may be instrumental in the execution of apoptosis following caspase activation. Mice that lack this gene have the ability to regenerate damaged or missing tissue. Multiple alternatively spliced variants have been found for this gene.




Western blot analysis of extracts of various cell lines using p21 Polyclonal Antibody at dilution of 1:1000.


Western blot analysis of extracts of 293T cells using p21 Polyclonal Antibody at dilution of 1:1000.



Immunofluorescence analysis of C6 cells using p21 Polyclonal Antibody at dilution of 1:100. Blue: DAPI for nuclear staining.



Immunofluorescence analysis of HeLa cells using p21 Polyclonal Antibody at dilution of 1:100. Blue: DAPI for nuclear staining.



Immunofluorescence analysis of NIH/3T3 cells using p21 Polyclonal Antibody at dilution of 1:100. Blue: DAPI for nuclear staining.