




Product name:	Phospho-Catenin- β (S33/S37/T41)
Cat number:	ABP-0524
Conjugate:	Unconjugated
Size:	100 ug
Clone:	Poly
Concentration:	1mg/ml
Host:	Rb
Isotype:	IgG
Immunogen:	A synthetic phosphorylated peptide around S33 & T41 of human beta Catenin
Reactivity:	Hu, Ms, Rt
Applications:	Western Blot: 1:500 - 1:2000 Immunohistochemistry: 1:50 - 1:200
Molecular Weight:	100kDa
Purification:	Affinity purification
Form:	liquid
Buffer:	PBS with 0.02% sodium azide, 50% glycerol, pH7.3.
Storage:	Store at -20°C. Avoid freeze / thaw cycles.
Background:	The protein encoded by this gene is part of a complex of proteins that constitute adherens junctions (AJs). AJs are necessary for the creation and maintenance of epithelial cell layers by regulating cell growth and adhesion between cells. The encoded protein also anchors the actin cytoskeleton and may be responsible for transmitting the contact inhibition signal that causes cells to stop dividing once the epithelial sheet is complete. Finally, this protein binds to the product of the APC gene, which is mutated in adenomatous polyposis of the colon. Mutations in this gene are a cause of colorectal cancer (CRC), pilomatixoma (PTR), medulloblastoma (MDB), and ovarian cancer. Alternative splicing results in multiple transcript variants.




Western blot analysis of extracts of NIH/3T3 cells, using Phospho- β -Catenin-S33/S37/T41 antibody at 1:1000 dilution. NIH/3T3 cells were treated by Calyculin A (10 μ l) at 37°C for 30 minutes after serum-starvation overnight. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. Lysates/proteins: 25 μ g per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL West Pico Plus. Exposure time: 1s.



Immunohistochemistry of paraffin embedded human colon carcinoma using Phospho- β -Catenin-S33/S37/T41 antibody at dilution of 1:100 (40x lens).



Immunohistochemistry of paraffin embedded rat skin using Phospho- β -Catenin-S33/S37/T41 antibody at dilution of 1:100 (40x lens).



Immunohistochemistry of paraffin embedded mouse kidney using Phospho- β -Catenin-S33/S37/T41 antibody at dilution of 1:100 (40x lens).