


Product name:	Phospho-ERK1/ERK2 (p44/p42 MAPK) (T202/Y204)
Cat number:	ABP-84133
Conjugate:	Unconjugated
Size:	100 ug
Clone:	Poly
Concentration:	1mg/ml
Host:	RB
Isotype:	IgG
Reactivity:	Hu, Ms, Rt
Applications:	WB 1:1000, IHC(P) 1:100, IP 1:50, IF-IC 1:100-500
Molecular Weight:	42, 44 kDa
Purification:	Polyclonal antibodies are produced by immunizing animals with a synthetic phosphopeptide corresponding to residues surrounding Thr202/Tyr204 of human p44 MAP kinase.
Form:	liquid
Buffer:	Supplied in 20 mM Tris-HCl, pH 8.0 10 mg/ml BSA ,0.05% Sodium Azide
Storage:	Store at -20°C. Do not aliquot the antibody.

Background:

Mitogen-activated protein kinases (MAPKs) are a widely conserved family of serine/threonine protein kinases involved in many cellular programs, such as cell proliferation, differentiation, motility, and death. The p44/42 MAPK (Erk1/2) signaling pathway can be activated in response to a diverse range of extracellular stimuli including mitogens, growth factors, and cytokines (1-3), and research investigators consider it an important target in the diagnosis and treatment of cancer (4). Upon stimulation, a sequential three-part protein kinase cascade is initiated, consisting of a MAP kinase kinase kinase (MAPKKK or MAP3K), a MAP kinase kinase (MAPKK or MAP2K), and a MAP kinase (MAPK). Multiple p44/42 MAP3Ks have been identified, including members of the Raf family, as well as Mos and Tpl2/COT. MEK1 and MEK2 are the primary MAPKKs in this pathway (5,6). MEK1 and MEK2 activate p44 and p42 through phosphorylation of activation loop residues Thr202/Tyr204 and Thr185/Tyr187, respectively. Several downstream targets of p44/42 have been identified, including p90RSK (7) and the transcription factor Elk-1 (8,9). p44/42 are negatively regulated by a family of dual-specificity (Thr/Tyr) MAPK phosphatases, known as DUSPs or MKPs (10), along with MEK inhibitors, such as U0126 and PD98059. Phospho-p44/42 MAPK (Erk1/2) (Thr202/Tyr204) Antibody detects endogenous levels of p44 and p42 MAP Kinase (Erk1 and Erk2) when phosphorylated either individually or dually at Thr202 and Tyr204 of Erk1 (Thr185 and Tyr187 of Erk2). The antibody does not cross-react with the corresponding phosphorylated residues of either JNK/ SAPK or p38 MAP Kinase, and does not cross-react with non-phosphorylated Erk1/2.



Specificity and sensitivity of Phospho-p44/42 MAPK (Erk1/2) (Thr202/Tyr204) Antibody. The antibody reacts specifically with as little as 0.25 ng of phosphorylated p42 MAP kinase and does not cross-react with up to 4 µg of nonphosphorylated p42 MAP kinase.



Anti - phospho-Erk 1,2 -Wash buffer: 1x Tris Buffered Saline (TBS); 0.1% Triton X-100 - Blocking buffer: 1xTBS; 0.1% Triton X-100; 5% BSA (used with the primary antibody) For western blots, incubate the membrane with antibody diluted in blocking buffer 2 hours at room temperature.