




Product name:	Phospho-GSK3 β (S9)
Cat number:	MAB-94624
Conjugate:	Unconjugated
Size:	100 ug
Clone:	5B3
Concentration:	1mg/ml
Host:	Rb
Isotype:	IgG
Reactivity:	Hu, Ms, Rt
Applications:	Western blotting 1:1000 ☐ Immunohistochemistry 1:100 ☐ Immunofluorescence (IF-IC) 1:200
Molecular Weight:	46 kDa
Purification:	Monoclonal antibody is produced by immunizing animals with a synthetic phosphopeptide corresponding to residues surrounding Ser9 of human GSK-3beta
Form:	liquid
Buffer:	Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 μ g/ml BSA, 50% glycerol and less than 0.02% sodium azide.
Storage:	Storage: at 4-8 $^{\circ}$ C for short term, at -20 $^{\circ}$ C for longer term.
Background:	Glycogen synthase kinase-3 (GSK-3) was initially identified as an enzyme that regulates glycogen synthesis in response to insulin (1). GSK-3 is a ubiquitously expressed serine/threonine protein kinase that phosphorylates and inactivates glycogen synthase. GSK-3 is a critical downstream element of the PI3 kinase/Akt cell survival pathway whose activity can be inhibited by Akt-mediated phosphorylation at Ser21 of GSK-3a and Ser9 of GSK-3beta (2,3). GSK-3 has been implicated in the regulation of cell fate in Dictyostelium and is a component of the Wnt signaling pathway required for Drosophila, Xenopus and mammalian development (4). GSK-3 has been shown to regulate cyclin D1 proteolysis and subcellular localization (5). Phospho-GSK-3beta (Ser9) (5B3) Rabbit mAb detects endogenous levels of GSK-3beta only when phosphorylated at Ser9. This antibody reacts with denatured components of bovine serum, including BSA.




Western blot analysis of extracts from PC-3 cells, untreated or LY294002/wortmannin-treated, using Phospho-GSK-3 β (Ser9) (Rabbit mAb (upper) or GSK-3 β Rabbit (lower).



Immunohistochemical analysis of paraffin embedded human breast carcinoma tissue using Phospho GSK3B - S9 antibody



Western blot analysis of extracts from GSK-3 α (-/-) (lanes 1,2) and GSK-3 β (-/-) (lanes 3,4) mouse embryonic fibroblast (MEF) cells, λ phosphatase or PDGF-treated, using Phospho-GSK-3 β (Ser9) (Rabbit mAb (upper) and α / β -Tubulin Antibody (lower).



Immunofluorescence staining of methanol-fixed HeLa Cells showing cytoplasmic staining using Phospho-GSK3B-S9 antibody