

|                          |                                                                                                  |
|--------------------------|--------------------------------------------------------------------------------------------------|
| <b>Product name:</b>     | POT1 Rabbit Polyclonal Antibody                                                                  |
| <b>Cat number:</b>       | ABN16380                                                                                         |
| <b>Conjugate:</b>        | Unconjugated                                                                                     |
| <b>Size:</b>             | 100μL                                                                                            |
| <b>Clone:</b>            | Polyclonal                                                                                       |
| <b>Concentration:</b>    | 1mg/ml                                                                                           |
| <b>Host:</b>             | Rabbit                                                                                           |
| <b>Isotype:</b>          | IgG                                                                                              |
| <b>Immunogen:</b>        | The antiserum was produced against synthesized peptide derived from human POT1. AA range:247-296 |
| <b>Reactivity:</b>       | Human,Monkey                                                                                     |
| <b>Applications:</b>     | WB 1:500-1:2000,IHC 1:50-1:300                                                                   |
| <b>Molecular Weight:</b> | 70kDa                                                                                            |
| <b>Purification:</b>     | Affinity purification                                                                            |
| <b>Form:</b>             | Liquid                                                                                           |
| <b>Buffer:</b>           | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.               |
| <b>Storage:</b>          | Store at 4°C short term. Aliquot and store at -20°C for 12 months. Avoid freeze/thaw cycles.     |

**Background:**

protection of telomeres 1(POT1) Homo sapiens This gene is a member of the telombin family and encodes a nuclear protein involved in telomere maintenance. Specifically, this protein functions as a member of a multi-protein complex that binds to the TTAGGG repeats of telomeres, regulating telomere length and protecting chromosome ends from illegitimate recombination, catastrophic chromosome instability, and abnormal chromosome segregation. Increased transcriptional expression of this gene is associated with stomach carcinogenesis and its progression. Alternatively spliced transcript variants have been described. [provided by RefSeq, Jul 2008],alternative products:A number of isoforms are produced,function:Component of the telomerase ribonucleoprotein (RNP) complex that is essential for the replication of chromosome termini. Is a component of the double-stranded telomeric DNA-binding TRF1 complex which is involved in the regulation of telomere length by cis-inhibition of telomerase. Also acts as a single-stranded telomeric DNA-binding protein and thus may act as a downstream effector of the TRF1 complex and may transduce information about telomere maintenance and/or length to the telomere terminus. Component of the shelterin complex (telosome) that is involved in the regulation of telomere length and protection. Shelterin associates with arrays of double-stranded TTAGGG repeats added by telomerase and protects chromosome ends; without its protective activity, telomeres are no longer hidden from the DNA damage surveillance and chromosome ends are inappropriately processed by DNA repair pathways. Binds to two or more telomeric single-stranded 5'-TTAGGG-3' repeats (G-strand) and with high specificity to a minimal telomeric single-stranded 5'-TAGGGTTAG-3' sequence. Binds telomeric single-stranded sequences internally or at proximity of a 3'-end. Its activity is TERT dependent but it does not increase TERT activity.,similarity:Belongs to the telombin family.,subcellular location:Colocalizes with telomeric DNA.,subunit:Homodimer or homooligomer. Component of the shelterin complex (telosome) composed of TERF1, TERF2, TINF2, TERF2IP, ACD and POT1. Binds single-stranded telomeric DNA as a monomer (By similarity). Component of the telomerase holoenzyme complex at least composed of TERT, DKC1, WDR79/TCAB1, NOP10, NHP2, GAR1, TEP1, EST1A, POT1 and a telomerase RNA template component (TERC). Found in a complex with TERF1, TINF2 and TNKS1. Interacts with TNKS1. Forms heterodimers with ACD.,tissue specificity:Ubiquitous.,