

Product name:	PRIC285 Rabbit Polyclonal Antibody
Cat number:	ABN16493
Conjugate:	Unconjugated
Size:	100µL
Clone:	Polyclonal
Concentration:	1mg/ml
Host:	Rabbit
Isotype:	IgG
Immunogen:	The antiserum was produced against synthesized peptide derived from human PRIC285. AA range:1909-1958
Reactivity:	Human,Rat,Mouse
Applications:	WB 1:500-1:2000,IHC 1:50-1:300,ELISA 1:2000-1:20000
Molecular Weight:	300kDa
Purification:	Affinity purification
Form:	Liquid
Buffer:	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.
Storage:	Store at 4°C short term. Aliquot and store at -20°C for 12 months. Avoid freeze/thaw cycles.
Background:	The protein encoded by this gene is a nuclear transcriptional co-activator for peroxisome proliferator activated receptor alpha. The encoded protein contains a zinc finger and is a helicase that appears to be part of the peroxisome proliferator activated receptor alpha interacting complex. This gene is a member of the DNA2/NAM7 helicase gene family. Alternatively spliced transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Jul 2008],caution:PubMed:12189208 experiments have been carried out partly in rat and partly in human.,domain:Contains 5 Leu-Xaa-Xaa-Leu-Leu (LXXLL) motifs. These motifs are not required for interaction with PPARG.,function:Helicase that acts as a transcriptional coactivator for a number of nuclear receptors including PPARA, PPARG, THRA, THRB and RXRA.,similarity:Belongs to the DNA2/NAM7 helicase family.,similarity:Contains 1 C2H2-type zinc finger.,subunit:Interacts with PPARA, ESR1, THRB and RXRA. It is probably part of the peroxisome proliferator activated receptor alpha interacting complex (PRIC). Also interacts with the DNA-binding domain of PPARG.,tissue specificity:Expressed in various tissues including heart, pancreas, skeletal muscle, colon, spleen, liver, kidney, lung, peripheral blood and placenta.,