

|                          |                                                                                                                                                                                              |
|--------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Product name:</b>     | SAM68 (1C18) Rabbit Monoclonal Antibody                                                                                                                                                      |
| <b>Cat number:</b>       | MABN17588                                                                                                                                                                                    |
| <b>Conjugate:</b>        | Unconjugated                                                                                                                                                                                 |
| <b>Size:</b>             | 100µL                                                                                                                                                                                        |
| <b>Clone:</b>            | Monoclonal                                                                                                                                                                                   |
| <b>Concentration:</b>    | 1mg/ml                                                                                                                                                                                       |
| <b>Host:</b>             | Rabbit                                                                                                                                                                                       |
| <b>Isotype:</b>          | IgG                                                                                                                                                                                          |
| <b>Immunogen:</b>        | A synthetic peptide of human SAM68                                                                                                                                                           |
| <b>Reactivity:</b>       | Human, Mouse                                                                                                                                                                                 |
| <b>Applications:</b>     | WB 1:1000-1:5000, IHC 1:200-1:500, ICC/IF 1:200-1:500, FC 1:100-1:1000                                                                                                                       |
| <b>Molecular Weight:</b> | 48kDa                                                                                                                                                                                        |
| <b>Purification:</b>     | Affinity purification                                                                                                                                                                        |
| <b>Form:</b>             | Liquid                                                                                                                                                                                       |
| <b>Buffer:</b>           | Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% New type preservative N and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle. |
| <b>Storage:</b>          | Store at 4°C short term. Aliquot and store at -20°C for 12 months. Avoid freeze/thaw cycles.                                                                                                 |

**Background:**

Recruited and tyrosine phosphorylated by several receptor systems, for example the T-cell, leptin and insulin receptors. Once phosphorylated, functions as an adapter protein in signal transduction cascades by binding to SH2 and SH3 domain-containing proteins. Recruited and tyrosine phosphorylated by several receptor systems, for example the T-cell, leptin and insulin receptors. Once phosphorylated, functions as an adapter protein in signal transduction cascades by binding to SH2 and SH3 domain-containing proteins. Role in G2-M progression in the cell cycle. Represses CBP-dependent transcriptional activation apparently by competing with other nuclear factors for binding to CBP. Also acts as a putative regulator of mRNA stability and/or translation rates and mediates mRNA nuclear export. Positively regulates the association of constitutive transport element (CTE)-containing mRNA with large polyribosomes and translation initiation. According to some authors, is not involved in the nucleocytoplasmic export of unspliced (CTE)-containing RNA species according to (PubMed:22253824). RNA-binding protein that plays a role in the regulation of alternative splicing and influences mRNA splice site selection and exon inclusion. Binds to RNA containing 5'-[AU]UAA- 3' as a bipartite motif spaced by more than 15 nucleotides. Binds poly(A). Can regulate CD44 alternative splicing in a Ras pathway- dependent manner (By similarity). In cooperation with HNRNPA1 modulates alternative splicing of BCL2L1 by promoting splicing toward isoform Bcl-X(S), and of SMN1 (PubMed:17371836, PubMed:20186123). Can regulate alternative splicing of NRXN1 and NRXN3 in the laminin G-like domain 6 containing the evolutionary conserved neuexin alternative spliced segment 4 (AS4) involved in neuexin selective targeting to postsynaptic partners. In a neuronal activity-dependent manner cooperates synergistically with KHDRBS2/SLIM-1 in regulation of NRXN1 exon skipping at AS4. The cooperation with KHDRBS2/SLIM-1 is antagonistic for regulation of NXRN3 alternative splicing at AS4 (By similarity).