Product name:	SCIN Rabbit Monoclonal Antibody
Cat number:	MABN02579
Conjugate:	Unconjugated
Size:	100µL
Clone:	Monoclonal
Concentration:	1mg/ml
Host:	Rabbit
Isotype:	IgG
Immunogen:	A synthetic peptide of human SCIN
Reactivity:	Human
Applications:	WB 1:500-1:1000
Molecular Weight:	Calculated MW: 80 kDa; Observed MW: 80 kDa
Purification:	Affinity Purification
Form:	Liquid
Buffer:	50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% Sodium azide and 0.05% BSA
Storage:	Store at 4°C short term. Aliquot and store at -20°C for 12 months. Avoid freeze/thaw cycles.
Background:	Ca2+-dependent actin filament-severing protein that has a regulatory function in exocytosis by affecting the organization of the microfilament network underneath the plasma membrane (PubMed:8547642, PubMed:26365202). Severing activity is inhibited by phosphatidylinositol 4,5-bis-phosphate (PIP2) . In vitro, also has barbed end capping and nucleating activities in the presence of Ca2+. Required for megakaryocyte differentiation, maturation, polyploidization and apoptosis with the release of platelet-like particles (PubMed:11568009). Plays a role in osteoclastogenesis (OCG) and actin cytoskeletal organization in osteoclasts . Regulates chondrocyte proliferation and differentiation . Inhibits cell proliferation and tumorigenesis. Signaling is mediated by MAPK, p38 and JNK pathways (PubMed:11568009).