

|                          |                                                                                                   |
|--------------------------|---------------------------------------------------------------------------------------------------|
| <b>Product name:</b>     | SF-1 Rabbit Polyclonal Antibody                                                                   |
| <b>Cat number:</b>       | ABN17780                                                                                          |
| <b>Conjugate:</b>        | Unconjugated                                                                                      |
| <b>Size:</b>             | 100µL                                                                                             |
| <b>Clone:</b>            | Polyclonal                                                                                        |
| <b>Concentration:</b>    | 1mg/ml                                                                                            |
| <b>Host:</b>             | Rabbit                                                                                            |
| <b>Isotype:</b>          | IgG                                                                                               |
| <b>Immunogen:</b>        | The antiserum was produced against synthesized peptide derived from human NR5A1. AA range:169-218 |
| <b>Reactivity:</b>       | Human,Mouse,Rat                                                                                   |
| <b>Applications:</b>     | WB 1:500-1:2000,IHC 1:50-1:300,ELISA 1:2000-1:20000                                               |
| <b>Molecular Weight:</b> | 50kDa                                                                                             |
| <b>Purification:</b>     | Affinity purification                                                                             |
| <b>Form:</b>             | Liquid                                                                                            |
| <b>Buffer:</b>           | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.                |
| <b>Storage:</b>          | Store at 4°C short term. Aliquot and store at -20°C for 12 months. Avoid freeze/thaw cycles.      |

**Background:**

The protein encoded by this gene is a transcriptional activator involved in sex determination. The encoded protein binds DNA as a monomer. Defects in this gene are a cause of XY sex reversal with or without adrenal failure as well as adrenocortical insufficiency without ovarian defect. [provided by RefSeq, Jul 2008],disease:Defects in NR5A1 are a cause of adrenocortical insufficiency without ovarian defect [MIM:184757]. The disease is characterized by severe 'slackness,' muscular hypotonia. There is decreased sodium, increased potassium and elevated ACTH.,disease:Defects in NR5A1 are a cause of XY sex reversal with or without adrenal failure [MIM:184757]. This disease is characterized by normal female external genitalia and retention of the uterus.,function:Transcriptional activator. Seems to be essential for sexual differentiation and formation of the primary steroidogenic tissues. Binds to the Ad4 site found in the promoter region of steroidogenic P-450 genes such as CYP11A, CYP11B and CYP21B. Also regulates the Muellerian inhibiting substance (AMH) gene as well as the AHCH and STAR genes. 5'-YCAAGGYC-3' and 5'-RRAGGTCA-3' are the consensus sequences for the recognition by NR5A1/FTZF1. The SFPQ-NONO-NR5A1/SF-1 complex binds to the CYP17 promoter and regulates basal and cAMP-dependent transcriptional activity. Binds phosphatidylcholine (By similarity). Binds phospholipids with a phosphatidylinositol (PI) headgroup, in particular PI(3,4)P2 and PI(3,4,5)P3.,PTM:Acetylation stimulates the transcriptional activity.,similarity:Belongs to the nuclear hormone receptor family.,similarity:Belongs to the nuclear hormone receptor family. NR5 subfamily.,similarity:Contains 1 nuclear receptor DNA-binding domain.,subunit:Binds DNA as a monomer. Interacts with NR0B2 and PPARGC1A (By similarity). Part of a complex consisting of SFPQ, NONO and NR5A1/SF-1. Interacts with NCOA2.,