

Product name:	SIP1 Rabbit Polyclonal Antibody
Cat number:	ABN17905
Conjugate:	Unconjugated
Size:	100µL
Clone:	Polyclonal
Concentration:	1mg/ml
Host:	Rabbit
Isotype:	IgG
Immunogen:	The antiserum was produced against synthesized peptide derived from human ZEB2. AA range:71-120
Reactivity:	Human,Mouse,Rat
Applications:	WB 1:500-1:2000,IHC 1:100-1:300,ICC/IF 1:50-1:200,ELISA 1:5000-1:20000
Molecular Weight:	157kDa
Purification:	Affinity purification
Form:	Liquid
Buffer:	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.
Storage:	Store at 4°C short term. Aliquot and store at -20°C for 12 months. Avoid freeze/thaw cycles.

Background:

The protein encoded by this gene is a member of the Zfh1 family of 2-handed zinc finger/homeodomain proteins. It is located in the nucleus and functions as a DNA-binding transcriptional repressor that interacts with activated SMADs. Mutations in this gene are associated with Hirschsprung disease/Mowat-Wilson syndrome. Alternatively spliced transcript variants have been found for this gene.[provided by RefSeq, Jan 2010],disease:Defects in ZEB2 are the cause of Hirschsprung disease-mental retardation syndrome (Hirschsprung disease) [MIM:235730]; also known as Mowat-Wilson syndrome (MWS). Hirschsprung disease is a rare autosomal dominant complex developmental disorder. Individuals with functional null mutations present with mental retardation, delayed motor development, epilepsy, and a wide spectrum of clinically heterogeneous features suggestive of neurocristopathies at the cephalic, cardiac, and vagal levels. Affected patients show an easily recognizable facial appearance with deep set eyes and hypertelorism, medially divergent, broad eyebrows, prominent columella, pointed chin and uplifted, notched ear lobes. Additionally, the phenotypic spectrum of facultative congenital anomalies includes short stature, microcephaly, Hirschsprung disease, malformations of the brain (agenesis of corpus callosum, cerebral atrophy) and eye (microphthalmia), seizures, congenital heart defects and genitourinary malformations, in particular hypospadias. The development of psychomotor skills and speech is delayed in most patients. Overall, the grade of mental retardation is at least moderate, but usually severe including characteristic abnormal behavior.,function:Transcriptional inhibitor that binds to DNA sequence 5'-CACCT-3' in different promoters. Represses transcription of E-cadherin.,PTM:Sumoylation on Lys-391 and Lys-866 is promoted by the E3 SUMO-protein ligase CBX4, and impairs interaction with CTBP1 and transcription repression activity.,similarity:Belongs to the delta-EF1/ZFH-1 C2H2-type zinc-finger family.,similarity:Contains 1 homeobox DNA-binding domain.,similarity:Contains 7 C2H2-type zinc fingers.,subunit:Binds activated SMAD1, activated SMAD2 and activated SMAD3; binding with SMAD4 is not detected (By similarity). Interacts with CBX4 and CTBP1.,