



Product name:	TAF15
Cat number:	AB-84381
Conjugate:	Unconjugated
Size:	100 ug
Clone:	POLY
Concentration:	1mg/ml
Host:	Rb
Isotype:	IgG
Immunogen:	Recombinant fusion protein containing a sequence corresponding to amino acids 1-280 of human TAF15
Reactivity:	Hu
Applications:	WB: 1:500 - 1:2000 IHC: 1:50 - 1:100 IF: 1:50 - 1:200 IP: 1:50 - 1:200
Molecular Weight:	80kDa
Purification:	Aff. Pur.
Form:	Liquid
Buffer:	Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH 7.3.
Storage:	Store at -20°C. Avoid freeze / thaw cycles.
Background:	This gene encodes a member of the TET family of RNA-binding proteins. The encoded protein plays a role in RNA polymerase II gene transcription as a component of a distinct subset of multi-subunit transcription initiation factor TFIID complexes. Translocations involving this gene play a role in acute leukemia and extraskeletal myxoid chondrosarcoma, and mutations in this gene may play a role in amyotrophic lateral sclerosis. Alternatively spliced transcript variants encoding multiple isoforms have been observed for this gene.



Western blot analysis of extracts of various cell lines, using TAF15 antibody at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL West Pico Plus. Exposure time: 5s.

Immunoprecipitation analysis of 200ug extracts of SKOV3 cells using 3ug TAF15 antibody. Western blot was performed from the immunoprecipitate using TAF15 antibody at a dilution of 1:1000.



Immunohistochemistry of paraffinembedded human liver cancer using TAF15 Rabbit pAb at dilution of 1:150 (40x lens).