

Product name:	TAT Rabbit Polyclonal Antibody
Cat number:	ABN18653
Conjugate:	Unconjugated
Size:	100µL
Clone:	Polyclonal
Concentration:	1mg/ml
Host:	Rabbit
Isotype:	IgG
Immunogen:	The antiserum was produced against synthesized peptide derived from human TAT. AA range:255-304
Reactivity:	Human,Mouse,Rat
Applications:	IHC 1:100-1:300,ICC/IF 1:50-1:200,ELISA 1:20000-1:40000
Purification:	Affinity purification
Form:	Liquid
Buffer:	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.
Storage:	Store at 4°C short term. Aliquot and store at -20°C for 12 months. Avoid freeze/thaw cycles.
Background:	This nuclear gene encodes a mitochondrial protein tyrosine aminotransferase which is present in the liver and catalyzes the conversion of L-tyrosine into p-hydroxyphenylpyruvate. Mutations in this gene cause tyrosinemia (type II, Richner-Hanhart syndrome), a disorder accompanied by major skin and corneal lesions, with possible mental retardation. A regulator gene for tyrosine aminotransferase is X-linked. [provided by RefSeq, Jul 2008],catalytic activity:L-tyrosine + 2-oxoglutarate = 4-hydroxyphenylpyruvate + L-glutamate.,cofactor:Pyridoxal phosphate.,disease:Defects in TAT are the cause of tyrosinemia type 2 (TYRO2) [MIM:276600]; also known as Richner-Hanhart syndrome. TYRO2 is an inborn error of metabolism characterized by elevations of tyrosine in the blood and urine, and oculocutaneous manifestations. Typical features include palmoplantar keratosis, painful corneal ulcers, and mental retardation.,pathway:Amino-acid degradation; L-phenylalanine degradation; acetoacetic acid and fumarate from L-phenylalanine: step 2/6.,similarity:Belongs to the class-I pyridoxal-phosphate-dependent aminotransferase family.,subunit:Homodimer.,