

Product name:	TRAF1 Rabbit Polyclonal Antibody
Cat number:	ABN19181
Conjugate:	Unconjugated
Size:	100µL
Clone:	Polyclonal
Concentration:	1mg/ml
Host:	Rabbit
Isotype:	IgG
Immunogen:	Synthesized peptide derived from TRAF1 at AA range: 191-240
Reactivity:	Human, Mouse
Applications:	WB 1:500-1:2000, ELISA 1:10000-1:20000
Molecular Weight:	46kDa
Purification:	Affinity purification
Form:	Liquid
Buffer:	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.
Storage:	Store at 4°C short term. Aliquot and store at -20°C for 12 months. Avoid freeze/thaw cycles.

Background:

TNF receptor associated factor 1 (TRAF1) Homo sapiens The protein encoded by this gene is a member of the TNF receptor (TNFR) associated factor (TRAF) protein family. TRAF proteins associate with, and mediate the signal transduction from various receptors of the TNFR superfamily. This protein and TRAF2 form a heterodimeric complex, which is required for TNF-alpha-mediated activation of MAPK8/JNK and NF-kappaB. The protein complex formed by this protein and TRAF2 also interacts with inhibitor-of-apoptosis proteins (IAPs), and thus mediates the anti-apoptotic signals from TNF receptors. The expression of this protein can be induced by Epstein-Barr virus (EBV). EBV infection membrane protein 1 (LMP1) is found to interact with this and other TRAF proteins; this interaction is thought to link LMP1-mediated B lymphocyte transformation to the signal transduction from TNFR family receptors. Three transcript variants encoding two different isoforms have domain: The coiled coil domain mediates homo- and hetero-oligomerization., domain: The MATH/TRAF domain binds to receptor cytoplasmic domains., function: Adapter protein and signal transducer that links members of the tumor necrosis factor receptor family to different signaling pathways by association with the receptor cytoplasmic domain and kinases. Mediates activation of NF-kappa-B and JNK and is involved in apoptosis. The TRAF1/TRAF2 complex recruits the apoptotic suppressors BIRC2 and BIRC3 to TNFRSF1B/TNFR2., similarity: Contains 1 MATH domain., subunit: Homotrimer (Probable). Heteromer with TRAF2 and associates with TNFRSF1B/TNFR2 through TRAF2. Associates with TNFRSF4, TNFRSF5/CD40, TNFRSF8/CD30, TNFRSF9/CD137, TNFRSF11A/RANK, TNFRSF18/AITR, TNFRSF17/BCMA, TNFRSF19/TROY, TNFRSF19L/RELT, XEDAR, EDAR, Epstein-Barr virus BNFL1/LMP-1, TANK/ITRAF, TRAIIP and RIPK2. Interacts with BIRC2 and BIRC3 N-terminus. Interacts with NFATC2IP and with HIVEP3.,