

|                          |                                                                                                   |
|--------------------------|---------------------------------------------------------------------------------------------------|
| <b>Product name:</b>     | TRAIL Rabbit Polyclonal Antibody                                                                  |
| <b>Cat number:</b>       | ABN19193                                                                                          |
| <b>Conjugate:</b>        | Unconjugated                                                                                      |
| <b>Size:</b>             | 100µL                                                                                             |
| <b>Clone:</b>            | Polyclonal                                                                                        |
| <b>Concentration:</b>    | 1mg/ml                                                                                            |
| <b>Host:</b>             | Rabbit                                                                                            |
| <b>Isotype:</b>          | IgG                                                                                               |
| <b>Immunogen:</b>        | The antiserum was produced against synthesized peptide derived from human TNFSF10. AA range:31-80 |
| <b>Reactivity:</b>       | Human,Rat,Mouse                                                                                   |
| <b>Applications:</b>     | WB 1:500-1:2000,IHC 1:100-1:300,ICC/IF 1:200-1:1000,ELISA 1:10000-1:20000                         |
| <b>Molecular Weight:</b> | 30kDa                                                                                             |
| <b>Purification:</b>     | Affinity purification                                                                             |
| <b>Form:</b>             | Liquid                                                                                            |
| <b>Buffer:</b>           | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.                |
| <b>Storage:</b>          | Store at 4°C short term. Aliquot and store at -20°C for 12 months. Avoid freeze/thaw cycles.      |

**Background:**

The protein encoded by this gene is a cytokine that belongs to the tumor necrosis factor (TNF) ligand family. This protein preferentially induces apoptosis in transformed and tumor cells, but does not appear to kill normal cells although it is expressed at a significant level in most normal tissues. This protein binds to several members of TNF receptor superfamily including TNFRSF10A/TRAILR1, TNFRSF10B/TRAILR2, TNFRSF10C/TRAILR3, TNFRSF10D/TRAILR4, and possibly also to TNFRSF11B/OPG. The activity of this protein may be modulated by binding to the decoy receptors TNFRSF10C/TRAILR3, TNFRSF10D/TRAILR4, and TNFRSF11B/OPG that cannot induce apoptosis. The binding of this protein to its receptors has been shown to trigger the activation of MAPK8/JNK, caspase 8, and caspase 3. Alternatively spliced transcript variants encoding different isoforms have been found for this gene. [provincofactor: Binds 1 zinc ion per trimer., function: Cytokine that binds to TNFRSF10A/TRAILR1, TNFRSF10B/TRAILR2, TNFRSF10C/TRAILR3, TNFRSF10D/TRAILR4 and possibly also to TNFRSF11B/OPG. Induces apoptosis. Its activity may be modulated by binding to the decoy receptors TNFRSF10C/TRAILR3, TNFRSF10D/TRAILR4 and TNFRSF11B/OPG that cannot induce apoptosis., similarity: Belongs to the tumor necrosis factor family., subunit: Homotrimer., tissue specificity: Widespread; most predominant in spleen, lung and prostate.,