




Product name:	Tyrosine Hydroxylase Rabbit Polyclonal Antibody
Cat number:	ABE3600
Conjugate:	unconjugated
Size:	100 ug
Clone:	Polyclonal
Concentration:	1 mg/ml
Host:	Rabbit
Isotype:	IgG
Immunogen:	The antiserum was produced against synthesized peptide derived from human Tyrosine Hydroxylase. AA range:41-90
Reactivity:	Human;Mouse;Rat
Applications:	Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/20000. Not yet tested in other applications.
Molecular Weight:	55kD
Purification:	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Form:	liquid
Buffer:	Liquid in PBS containing 50% glycerol, 0.5%BSAand0.02% sodium azide
Storage:	-20°C/1 year
Background:	The protein encoded by this gene is involved in the conversion of tyrosine to dopamine. It is the rate-limiting enzyme in the synthesis of catecholamines, hence plays a key role in the physiology of adrenergic neurons. Mutations in this gene have been associated with autosomal recessive Segawa syndrome. Alternatively spliced transcript variants encoding different isoforms have been noted for this gene. [provided by RefSeq, Jul 2008],




Western Blot analysis of various cells using
TH Polyclonal Antibody



Immunohistochemical analysis of paraffin-embedded Human brain.
Antibody was diluted at 1:100(4° overnight). High-pressure and
temperature Tris-EDTA,pH8.0 was used for antigen retrieval.
Negative contrl (right) obtained from antibody was pre-absorbed by
i



Immunofluorescence analysis of HUVEC cells, using Tyrosine
Hydroxylase Antibody. The picture on the right is blocked with the
synthesized peptide.



Immunohistochemistry analysis of paraffin-embedded human brain
tissue, using Tyrosine Hydroxylase Antibody. The picture on the
right is blocked with the synthesized peptide.