




Product name:	YAP
Cat number:	MAB-94469
Size:	100µg
Clone:	D8H1X
Concentration:	1mg/ml
Host:	Rb
Isotype:	IgG
Reactivity:	Hu,Ms,Rt
Applications:	Western blotting 1:1000 Immunoprecipitation 1:20-1:50 Immunohistochemistry (Paraffin) Unmasking buffer: Immunofluorescence IF Size: 100 ug Concentration: 1mg/ml Clone: D8H1X 1:50-1:200 1:50-1:200
Molecular Weight:	65-75 kDa
Purification:	Monoclonal antibody is produced by immunizing animals with recombinant protein specific to the carboxy terminus of human YAP protein. The epitope corresponds to a region surrounding Pro435 of human YAP isoform 1. This sequence region is 100% conserved among all known isoforms of human YAP protein.
Form:	Liquid
Buffer:	PBS with 0.02%sodium azide, 50% glycerol, pH7.3.
Background:	YAP (Yes-associated protein, YAP65) was identified based on its ability to associate with the SH3 domain of Yes. It also binds to other SH3 domain-containing proteins such as Nck, Crk, Src, and Abl (1). In addition to the SH3 binding motif, YAP contains a PDZ interaction motif, a coiled-coil domain, and WW domains (2-4). While initial studies of YAP all pointed towards a role in anchoring and targeting to specific subcellular compartments, subsequent studies showed that YAP is a transcriptional coactivator by virtue of its WW domain interacting with the PY motif (PPxY) of the transcription factor PEBP2 and other transcription factors (5,6). In its capacity as a transcriptional coactivator, YAP is now widely recognized as a central mediator of the Hippo Pathway, which plays a fundamental and widely conserved role in regulating tissue growth and organ size. Upon phosphorylation at Ser127 by LATS1/2 kinases, YAP translocates to the cytoplasm, where it is sequestered through association with 14-3-3 proteins in an Akt-dependent manner (6-8).YAP (D8H1X) XP® Rabbit mAb recognizes endogenous levels of total YAP protein.




Western blot analysis of extracts of various cell lines, using YAP1 antibody at 1:1000 dilution. Secondary antibody: HRP Goat Anti- Rabbit IgG (H+L) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL West Pico Plus Exposure time: 10s.




Western blot analysis of extracts from normal (control) and YAP1 knockout (KO) HeLa cells, using YAP1 antibody at 1:1000 dilution. Secondary antibody: HRP Goat Anti- Rabbit IgG (H+L) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 90s.



Immunohistochemistry of paraffinembedded human placenta using YAP1 antibody at dilution of 1:100 (40x lens).



Immunofluorescence analysis of C6 cells using YAP1 antibody at dilution of 1:100. Blue: DAPI for nuclear staining.



Immunoprecipitation analysis of 200ug extracts of HeLa cells, using 3 ug YAP1 antibody. Western blot was performed from the immunoprecipitate using YAP1 antibody (at a dilution of 1:1000).