




Cat. No:	ABP10037
Conjugate:	Unconjugated
Size:	100 ug
Clone:	Poly
Concentration:	1mg/ml
Host:	Rabbit
Isotype:	IgG
Immunogen:	Synthesized peptide derived from human Akt around the phosphorylation site of T308.
Reactivity:	Human, Mouse, Rat
Applications:	Western Blot: 1:500-1:2000 Immunofluorescence: 1:50-200 Immunohistochemistry: 1:100-1:300 ELISA: 1:20000
Molecular Weight:	56 kDa
Purification:	The antibody was affinity-purified from rabbit antiserum by affinity chromatography using epitope-specific immunogen
Synonyms:	Anti-Rac-Alpha Serine/Threonine-Protein Kinase antibody Anti-Protein Kinase B antibody Anti-Pkb antibody Anti-Protein Kinase B Alpha antibody Anti-Pkb Alpha antibody Anti-Proto-Oncogene C-Akt antibody Anti- Rac-Pk-Alpha antibody Anti-AKT1 PKB RAC antibody
Background:	<p>Akt is a protein kinase consisting of three proteins (Akt1, Akt2 and Akt3) encoded by the AKT gene which is approximately 55 kDa. Akt is localized to the cytoplasm, nucleus and cell membrane and is involved pathways such as RET signalling, regulation of lipid metabolism, insulin signalling, generic cascades, common cytokine receptor gamma-chain family signalling pathways and apoptotic pathways. Akt regulates proliferation, cell survival, growth, angiogenesis and tumor formation. This is mediated through serine and/or threonine phosphorylation of a range of downstream substrates. Akt1 and Akt2 are expressed in all human cell types and AKT 3 is highly expressed in brain, lung and kidney. Mutations in the AKT1 gene can result in breast cancer and colorectal cancer. AKT was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. This primary antibody specifically binds to endogenous Akt protein which only binds about T308 when T308 is phosphorylated. Phospho-Akt (T308) polyclonal antibody detects endogenous levels of Akt protein only when phosphorylated at T308. Tissue Specificity: Expressed in prostate cancer and levels increase from the normal to the malignant state (at protein level). Expressed in all human cell types so far analyzed. The Tyr-176 phosphorylated form shows a significant increase in expression in breast cancers during the progressive stages i.e. normal to hyperplasia (ADH), ductal carcinoma in situ (DCIS), invasive ductal carcinoma (IDC) and lymph node metastatic (LNMM) stages.</p>
Form:	liquid
Buffer:	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Storage:	Store at -20°C, and avoid repeat freeze-thaw cycles.




Western blot (WB) analysis of AKT (phospho Thr308) polyclonal antibody.




Western blot analysis of lysates from A431 and HeLa cells treated with UV or insulin 0.01U/ml, using AKT p-308 Antibody. Primary antibody was diluted at 1:1000 4° over night, secondary antibody was diluted at 1:10000, 37° 1hour.




Immunohistochemical analysis of human-appendix tissue. Anti-Phospho-AKT (T308) at 1:200 (4°C, overnight). Antigen retrieval - Sodium Citrate pH6 (>98°C, 20min).




Immunohistochemical analysis of mouse spleen tissue. Anti-Phospho-AKT (T308) at 1:200 (4°C, overnight). Antigen retrieval - Sodium Citrate pH6 (>98°C, 20min).



Immunohistochemical analysis of mouse lung tissue. Anti-Phospho-AKT (T308) at 1:200 (4°C, overnight). Antigen retrieval - Sodium Citrate pH6 (>98°C, 20min).



Immunohistochemical analysis of rat spinal cord tissue. was diluted at 1:200 (4°C, overnight). Sodium citrate pH6.0 was used for antibody retrieval (>98°C, 20min).



Immunofluorescence analysis of human-lung tissue.
1, Akt (phospho Thr308) Polyclonal Antibody (red) was diluted at 1:200 (4°C, overnight).
2, Cy3 labeled Secondary antibody was diluted at 1:300 (room temperature, 50min).
3, Picture B: DAPI (blue) 10min.
Picture A: Target.
Picture B: DAPI.
Picture C: merge of A+B

**For Research use only
IMMUNOLOGICAL SCIENCES**