

IS-22412 Goat Anti-Rat IgG (H+L) HRP Conjugated Secondary Antibodies


Size: **500 ul**

Product Description: This product is prepared by labeling IgG (H+L) or recombinant streptavidin with horseradish peroxidase. HRP conjugates can be used for western blot, ELISA, immunohistochemistry and other standard immunoassay applications.


Applications: **Western Blot:** 1:5000 - 1:10000
Immunohistochemistry(paraffin-embedded tissues): 1:500 - 1:2000

Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.

Storage: Store at -20°C. Product is stable for at least 6 months from date of receipt when stored as recommended.



Western blot analysis of lysates from Rat IgG, using HRP Goat Anti-Rat IgG (H+L) at 1:1000 dilution. Lysates/proteins: 25 ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL West Pico Plus, Exposure time: 10s.



Western blot analysis of lysates from Rat IgG using HRP Goat Anti-Rat IgG (H+L) at 1:5000 dilution. Lysates/proteins: 50ng - 100ng per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL West Pico Plus. Exposure time: 30s.

**For Research use only
IMMUNOLOGICAL SCIENCES**