

IS20413 AP Goat Anti-Rabbit IgG (H+L)**Size:** 500 ul**Source:** Goat**Isotype:** Alkaline Phosphatase conjugated IgG**Purification:** Affinity purification**Immunogen:** Rabbit IgG

Background: Secondary antibodies are affinity-purified antibodies which will work with target-specific primary antibody in the detection, sorting or purification of its specified target. Secondary antibodies offer increased versatility enabling users to use many detection systems (e.g. HRP, AP, fluorescence). They can also provide greater sensitivity through signal amplification as multiple secondary antibodies. Most commonly, secondary antibodies are generated by immunizing the host animal (different from host species of primary antibody) with a pooled population of normal immunoglobulins from the host species of primary antibody and can be further purified and modified (i.e. antibody fragmentation, label conjugation, etc.) to ensure well-characterized specificity to corresponding normal immunoglobulins.

Applications: Western Blot: 1:5000 - 1:50000**Buffer:** PBS with 0.025% Sodium Azide, 0.75% BSA, 50% glycerol, pH 7.3**Storage:** Store at -20°C. Avoid freeze / thaw cycles**References: References for AP Goat Anti-Rabbit IgG (H+L)**

Product: AP Goat Anti-Rabbit IgG (H+L)

Journal: The Journal of steroid biochemistry and molecular biology

Application: WB

IF: 4.292

Species: Homo sapiens

PMID: [31376461](https://pubmed.ncbi.nlm.nih.gov/31376461/)

Title: Estrogen regulates forkhead transcription factor 2 to promote apoptosis of human ovarian granulosa-like tumor cells.