




<b>Product name:</b>	β-Tubulin Monoclonal Antibody(5G3)
<b>Cat number:</b>	MABN80021
<b>Conjugate:</b>	unconjugated
<b>Size:</b>	100 μL
<b>Clone:</b>	5G3
<b>Concentration:</b>	1mg/ml
<b>Host:</b>	Mouse
<b>Immunogen:</b>	Synthetic Peptide of β-Tubulin
<b>Applications:</b>	WB 1:5000, IHC-P 1:200, IF-P/IF-F/IF-ICC 1:200
<b>Purification:</b>	Affinity purification
<b>Form:</b>	liquid
<b>Buffer:</b>	Liquid in PBS,pH 7.4, containing 50% glycerol2% BSA and New type preservative N.
<b>Storage:</b>	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
<b>Source:</b>	Mouse


**For Research Use Only**

**IMMUNOLOGICAL SCIENCES**


Web-site: <https://immunologicalsciences.com> - E-mail: [info@immunologicalsciences.com](mailto:info@immunologicalsciences.com)



Western blot analysis of A549 (1) , Rat brain (2) ,Mouse brain (3) ,Chicken lung (4) and Rabbit testis (5),Sheep muscle (6) with  $\beta$ -tubulin mouse mAb(5G3) diluted at 1:5000.



IHC Staining of Human colon tissue with  $\beta$ -tubulin mouse mAb(5G3) diluted at 1:200.



IF analysis of HeLa with  $\beta$ -Tubulin Monoclonal Antibody(Left) and DAPI (Right) diluted at 1:100.