
Product name:	GFAP(5C8)Mouse Monoclonal Antibody
Cat number:	MABN11411
Conjugate:	Unconjugated
Size:	100ul
Clone:	5C8
Concentration:	1mg/ml
Host:	Mouse
Isotype:	IgG
Immunogen:	Synthetic Peptide of GFAP
Reactivity:	Human,Rat,Mouse
Applications:	WB 1:2000-1:5000,IHC 1:50-1:300,ICC/IF 1:100-1:200
Molecular Weight:	45kDa
Purification:	Affinity purification
Form:	liquid
Buffer:	PBS, pH 7.4, containing 0.5%protective protein, 0.02% New type preservative N as Preservative and 50% Glycerol.
Storage:	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
Synonyms:	GFAP; Glial fibrillary acidic protein; GFAP
Source:	Mouse
Background:	<p>This gene encodes one of the major intermediate filament proteins of mature astrocytes. It is used as a marker to distinguish astrocytes from other glial cells during development. Mutations in this gene cause Alexander disease, a rare disorder of astrocytes in the central nervous system. Alternative splicing results in multiple transcript variants encoding distinct isoforms. [provided by RefSeq, Oct 2008],alternative products:Isoforms differ in the C-terminal region which is encoded by alternative exons,disease:Defects in GFAP are a cause of Alexander disease (ALEXD) [MIM:203450]. Alexander disease is a rare disorder of the central nervous system. It is a progressive leukoencephalopathy whose hallmark is the widespread accumulation of Rosenthal fibers which are cytoplasmic inclusions in astrocytes. The most common form affects infants and young children, and is characterized by progressive failure of central myelination, usually leading to death usually within the first decade. Infants with Alexander disease develop a leukoencephalopathy with macrocephaly, seizures, and psychomotor retardation. Patients with juvenile or adult forms typically experience ataxia, bulbar signs and spasticity, and a more slowly progressive course.,function:GFAP, a</p>

For Research Use Only

IMMUNOLOGICAL SCIENCES




class-III intermediate filament, is a cell-specific marker that, during the development of the central nervous system, distinguishes astrocytes from other glial cells.,online information:GFAP entry,similarity:Belongs to the intermediate filament family.,subcellular location:Associated with intermediate filaments.,subunit:Interacts with SYNM (By similarity). Isoform 3 interacts with PSEN1 (via N-terminus),tissue specificity:Expressed in cells lacking fibronectin.,


For Research Use Only

IMMUNOLOGICAL SCIENCES


Web-site: <https://immunologicalsciences.com> - E-mail: info@immunologicalsciences.com




Immunohistochemical analysis of paraffin-embedded Human-liver tissue.
1,GFAP Monoclonal Antibody5C8 was diluted at 1:2004°C,overnight.
2, Sodium citrate pH 6.0 was used for antibody retrieval>98°C,20min.
3,Secondary antibody was diluted at 1:200room tempeRature, 30min.
Negative control was used by secondary antibody only.




Immunohistochemical analysis of paraffin-embedded Rat-heart tissue.
1,GFAP Monoclonal Antibody5C8 was diluted at 1:2004°C,overnight.
2, Sodium citrate pH 6.0 was used for antibody retrieval>98°C,20min.
3,Secondary antibody was diluted at 1:200room tempeRature, 30min.
Negative control was used by secondary antibody only.




Immunohistochemical analysis of paraffin-embedded Mouse-kidney tissue. 1,GFAP Monoclonal Antibody5C8 was diluted at 1:2004°C,overnight. 2, Sodium citrate pH 6.0 was used for antibody retrieval>98°C,20min. 3,Secondary antibody was diluted at 1:200room tempeRature, 30min. Negative control was used by secondary antibody only.



Immunofluorescence analysis of Mouse-brain tissue. 1,GFAP Monoclonal Antibody5C8red was diluted at 1:2004°C,overnight. 2, Cy3 labeled Secondary antibody was diluted at 1:300room temperature, 50min.3, Picture B: DAPIblue 10min. Picture A:Target. Picture B: DAPI. Picture C: merge of A+B



Immunofluorescence analysis of Rat-brain tissue. 1,GFAP Monoclonal Antibody5C8red was diluted at 1:2004°C,overnight. 2, Cy3 labeled Secondary antibody was diluted at 1:300room temperature, 50min.3, Picture B: DAPIblue 10min. Picture A:Target. Picture B: DAPI. Picture C: merge of A+B



Western blot analysis of Rat Brain Tissue, diluted at 1:5000.

**For Research Use Only
IMMUNOLOGICAL SCIENCES**