




Product name:	NMDAR1 (5R10) Rabbit Monoclonal Antibody
Cat number:	MABN14756
Conjugate:	Unconjugated
Size:	100ul
Clone:	5R10
Concentration:	1mg/ml
Host:	Rabbit
Isotype:	IgG
Immunogen:	A synthetic peptide of human NMDAR1
Reactivity:	Human,Mouse,Rat
Applications:	WB 1:1000-1:2000,ICC/IF 1:20-1:50
Molecular Weight:	105kDa
Purification:	Affinity purification
Form:	liquid
Buffer:	Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% New type preservative N and 0.05% protective protein.
Storage:	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
Synonyms:	GluN1; NMD-R1; GRIN1; NMDAR1
Source:	Rabbit
Background:	<p>N-methyl-D-aspartate receptor (NMDAR) forms a heterodimer of at least one NR1 and one NR2A-D subunit. Multiple receptor isoforms with distinct brain distributions and functional properties arise by selective splicing of the NR1 transcripts and differential expression of the NR2 subunits. Component of NMDA receptor complexes that function as heterotetrameric, ligand-gated ion channels with high calcium permeability and voltage-dependent sensitivity to magnesium. Channel activation requires binding of the neurotransmitter glutamate to the epsilon subunit, glycine binding to the zeta subunit, plus membrane depolarization to eliminate channel inhibition by Mg(2+) (PubMed:7685113, PubMed:28126851, PubMed:26919761, PubMed:26875626, PubMed:28105280). Sensitivity to glutamate and channel kinetics depend on the subunit composition (PubMed:26919761).</p>

For Research Use Only

IMMUNOLOGICAL SCIENCES

Web-site: <https://immunologicalsciences.com> - E-mail: info@immunologicalsciences.com



Western blot detection of NMDAR1 in
Rat brain, Jurkat cell lysates using
NMDAR1 antibody (1:1000 diluted).