




---

<b>Product name:</b>	CD338 Mouse Monoclonal Antibody
<b>Cat number:</b>	MABN82605
<b>Conjugate:</b>	Unconjugated
<b>Size:</b>	100ul
<b>Concentration:</b>	1mg/ml
<b>Host:</b>	Mouse
<b>Isotype:</b>	Mouse IgG2b
<b>Immunogen:</b>	Purified recombinant fragment of human CD338 (AA: extra mix) expressed in E. Coli.
<b>Reactivity:</b>	Human, Mouse
<b>Applications:</b>	WB 1:500-1:2000,IHC 1:200-1:1000,ICC 1:200-1:1000,ELISA 1:5000-1:20000,FC 1:200-1:400
<b>Molecular Weight:</b>	75kDa
<b>Purification:</b>	Affinity purification
<b>Form:</b>	liquid
<b>Buffer:</b>	Purified antibody in PBS with 0.05% sodium azide
<b>Storage:</b>	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
<b>Synonyms:</b>	MRX; MXR; ABCP; BCRP; BMDP; MXR1; ABC15; BCRP1; ABCG2; GOUT1; MXR-1; CDw338; UAQTL1; EST157481
<b>Source:</b>	Mouse
<b>Background:</b>	<p>The membrane-associated protein encoded by this gene is included in the superfamily of ATP-binding cassette (ABC) transporters. ABC proteins transport various molecules across extra- and intra-cellular membranes. ABC genes are divided into seven distinct subfamilies (ABC1, MDR/TAP, MRP, ALD, OABP, GCN20, White). This protein is a member of the White subfamily. Alternatively referred to as a breast cancer resistance protein, this protein functions as a xenobiotic transporter which may play a major role in multi-drug resistance. It likely serves as a cellular defense mechanism in response to mitoxantrone and anthracycline exposure. Significant expression of this protein has been observed in the placenta, which may suggest a potential role for this molecule in placenta tissue. Multiple transcript variants encoding different isoforms have been found for this gene.</p>


**For Research Use Only**

**IMMUNOLOGICAL SCIENCES**


Web-site: <https://immunologicalsciences.com> - E-mail: [info@immunologicalsciences.com](mailto:info@immunologicalsciences.com)




Black line: Control Antigen (100 ng); Purple line: Antigen (10ng); Blue line: Antigen (50 ng); Red line: Antigen (100 ng)




Western blot analysis using CD338 mouse mAb against A549 (1), HePG2 (2), Hela (3), and NIH/3T3 (4) cell lysate.



Immunofluorescence analysis of Hela cells using CD338 mouse mAb (green). Blue: DRAQ5 fluorescent DNA dye. Red: Actin filaments have been labeled with Alexa Fluor- 555 phalloidin.



Flow cytometric analysis of Hela cells using CD338 mouse mAb (green) and negative control (red).



Flow cytometric analysis of HePG2 cells using CD338 mouse mAb (green) and negative control (red).

**For Research Use Only**

**IMMUNOLOGICAL SCIENCES**