



Product name:	Glut4 Rabbit Polyclonal Antibody
Cat number:	ABN11504
Conjugate:	Unconjugated
Size:	100 ug
Concentration:	1mg/ml
Host:	Rabbit
Isotype:	IgG
Immunogen:	The antiserum was produced against synthesized peptide derived from the N-terminal region of human SLC2A4. AA range:21-70
Reactivity:	Human,Mouse,Rat
Applications:	WB 1:500-1:2000,IHC 1:100-1:300,ICC/IF 1:100-1:300,ELISA 1:10000-1:20000
Molecular Weight:	56kDa
Purification:	Affinity purification
Form:	Liquid
Buffer:	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
Storage:	Aliquot and store at -20°C (valid for 12 months). Avoid freeze/thaw cycles.
Synonyms:	SLC2A4; GLUT4; Solute carrier family 2, facilitated glucose transporter member 4; Glucose transporter type 4, insulin-responsive; GLUT-4
Source:	Rabbit
Background:	<p>This gene is a member of the solute carrier family 2 (facilitated glucose transporter) family and encodes a protein that functions as an insulin-regulated facilitative glucose transporter. In the absence of insulin, this integral membrane protein is sequestered within the cells of muscle and adipose tissue. Within minutes of insulin stimulation, the protein moves to the cell surface and begins to transport glucose across the cell membrane. Mutations in this gene have been associated with noninsulin-dependent diabetes mellitus (NIDDM). [provided by RefSeq, Jul 2008],disease:Defects in SLC2A4 may be a cause of noninsulin-dependent diabetes mellitus (NIDDM) [MIM:125853]. Defects in SLC2A4 may be a cause of certain post-receptor defects in NIDDM. The variant in position Ile-383 is found in a small number of NIDDM patients, but seems not to be found in nondiabetic subjects.,function:Insulin-regulated facilitative glucose transporter.,miscellaneous:Insulin-stimulated phosphorylation of TBC1D4 is required for GLUT4 translocation.,online information:GLUT4 entry,PTM:Sumoylated.,similarity:Belongs to the major facilitator superfamily. Sugar transporter (TC 2.A.1.1) family. Glucose transporter</p>

For Research Use Only

IMMUNOLOGICAL SCIENCES

Web-site: <https://immunologicalsciences.com> - E-mail: info@immunologicalsciences.com




subfamily.,subcellular location:Localizes primarily to the perinuclear region, undergoing continued recycling to the plasma membrane where it is rapidly reinternalized. The dileucine internalization motif is critical for intracellular sequestration.,subunit:Binds to DAXX. Interacts via its N-terminus with SRFBP1.,tissue specificity:Skeletal and cardiac muscles; brown and white fat.


For Research Use Only

IMMUNOLOGICAL SCIENCES


Web-site: <https://immunologicalsciences.com> - E-mail: info@immunologicalsciences.com




Immunohistochemical analysis of paraffin-embedded Rat-lung tissue. 1, Glut4 Polyclonal Antibody was diluted at 1:2004°C, overnight. 2, Sodium citrate pH 6.0 was used for antibody retrieval >98°C, 20min. 3, Secondary antibody was diluted at 1:200 room temperature, 30min. Negative control was used by secondary antibody only.




Immunohistochemical analysis of paraffin-embedded Rat-kidney tissue. 1, Glut4 Polyclonal Antibody was diluted at 1:2004°C, overnight. 2, Sodium citrate pH 6.0 was used for antibody retrieval >98°C, 20min. 3, Secondary antibody was diluted at 1:200 room temperature, 30min. Negative control was used by secondary antibody only.




Immunohistochemical analysis of paraffin-embedded Rat-spleen tissue. 1, Glut4 Polyclonal Antibody was diluted at 1:2004°C, overnight. 2, Sodium citrate pH 6.0 was used for antibody retrieval >98°C, 20min. 3, Secondary antibody was diluted at 1:200 room temperature, 30min. Negative control was used by secondary antibody only.



Immunohistochemical analysis of paraffin-embedded Mouse-heart tissue. 1, Glut4 Polyclonal Antibody was diluted at 1:2004°C, overnight. 2, Sodium citrate pH 6.0 was used for antibody retrieval >98°C, 20min. 3, Secondary antibody was diluted at 1:200 room temperature, 30min. Negative control was used by secondary antibody only.




Immunohistochemical analysis of paraffin-embedded Mouse-liver tissue. 1, Glut4 Polyclonal Antibody was diluted at 1:2004°C, overnight. 2, Sodium citrate pH 6.0 was used for antibody retrieval >98°C, 20min. 3, Secondary antibody was diluted at 1:200 room temperature, 30min. Negative control was used by secondary antibody only.



Immunohistochemical analysis of paraffin-embedded Mouse-kidney tissue. 1, Glut4 Polyclonal Antibody was diluted at 1:2004°C, overnight. 2, Sodium citrate pH 6.0 was used for antibody retrieval >98°C, 20min. 3, Secondary antibody was diluted at 1:200 room temperature, 30min. Negative control was used by secondary antibody only.

For Research Use Only

IMMUNOLOGICAL SCIENCES



Western Blot analysis of SKOV3 cells
using Glut4 Polyclonal Antibody..
Secondary antibody was diluted at
1:20000

**For Research Use Only
IMMUNOLOGICAL SCIENCES**

Web-site: <https://immunologicalsciences.com> - E-mail: info@immunologicalsciences.com